arXiv:hep-th/9402013v1 3 Feb 1994

RBI-TH-2/94
PMF-TH-1/94
January 1994

MAPPINGS OF MULTIMODE BOSE ALGEBRA PRESERVING NUMBER OPERATORS

M. Dorešić[1] , S.Meljanac[1] and M.Mileković[2,+]

[1] *Rudjer Bošković Institute, Bijenička c.54, 41001 Zagreb, Croatia*

[2] *Prirodoslovno-Matematički Fakultet, Department of Theoretical Physics*
Bijenička c.32, 41000 Zagreb, Croatia

[+] *e-mail: marijan@phy.hr*

Abstract

We define a class of deformed multimode oscillator algebras which possess number operators and can be mapped to multimode Bose algebra. We construct and discuss the states in the Fock space and the corresponding number operators.

Recently, much interest has ben devoted to the study of quantum groups [1] and to generalizations (deformations) of oscillator algebras [2][3][4][5][6]. Any type of single-mode deformed oscillator can be related (mapped) to a single-mode Bose oscillator . Deformations of multimode oscillators have also been studied [7]. For Pusz-Woronowitz oscillators covariant under the $SU_q(n)(SU_q(n|m))$ quantum algebra (superalgebra) [2][7] and anyonic-type algebras [8] there exist mappings to multimode Bose algebra. However, there are deformed generalized quon algebras [9] for which such mappings do not exist. It is essential that all these algebras are associative and the R-matrix approach to them has been pursued [10]. Multimode deformed oscillator algebras are important in possible physical applications, for example in q-deformed field theory [8] and generalized statistics [12].

Our aim is to define and analyze a general class of deformed multimode oscillator algebras which possess well-defined number operators for each type of oscillator, and can be mapped to a multimode Bose algebra. These mappings can be viewed as generalized Jordan-Wigner/Klein transformations. We discuss the corresponding algebras. For special values of parameters, one recovers all known such algebras [2][3] [5][6][7][8][10]. Finally, we construct and discuss the states in the Fock space and the corresponding number operators for deformed oscillators.

Let us define the annihilation and creation operators b_i, b_i^+, $(i \in S)$ satisfying the Bose algebra

$$\begin{gathered} [b_i, b_j^+] = \delta_{ij}, \quad \forall i, j \in S \\ [b_i, b_j] = 0 \end{gathered} \tag{1}$$

where $S = 1, 2, ...n$ or S is a set of sites on a lattice. The number operators N_i

satisfy

$$\begin{aligned} [N_i, b_j] &= -b_i \delta_{ij}, \quad \forall i, j \in S \\ [N_i, b_j^+] &= b_i^+ \delta_{ij} \\ N_i &= b_i^+ b_i, \quad \forall i, j \in S. \end{aligned} \tag{2}$$

Now we define generalized Jordan-Wigner transformations of the above Bose algebra, equations (1) and (2), as

$$a_i = b_i e^{\sum_j c_{ij} N_j} \sqrt{\frac{\varphi_i(N_i)}{N_i}} \tag{3}$$

where c_{ij} are complex numbers and $\varphi_i(N_i)$ is an arbitrary (complex) function with $\varphi_i(0) = 0$, $\lim_{\epsilon \to 0} \frac{\varphi_i(\epsilon)}{\epsilon} = 1$, $|\varphi_i(1)| = 1, \forall i \in S$. We assume that $|\varphi_i(N)|$ are bijective, monotonically increasing functions or that $\varphi_i(N) = \frac{1-(-)^N}{2}$, implying $a_i^2 = 0$. The mappings (3) generalize the mappings considered in [7][10].

It is important to note that the number operators are preserved, i.e.

$$N_i^{(a)} = N_i^{(b)} \equiv N_i, \quad \forall i \in S. \tag{4}$$

Then it is easy to find the corresponding deformed algebra:

$$\begin{aligned} a_i a_j &= e^{c_{ji} - c_{ij}} a_j a_i \quad i \neq j \\ a_i a_j^+ &= e^{c_{ij} + c_{ji}^*} a_j^+ a_i, \quad i \neq j \\ a_i a_i^+ &= |\varphi_i(N_i + 1)| e^{\sum_j (c_{ij} + c_{ij}^*) N_j} e^{(c_{ii} + c_{ii}^*)} \\ a_i^+ a_i &= |\varphi_i(N_i)| e^{\sum_j (c_{ij} + c_{ij}^*) N_j}. \end{aligned} \tag{5}$$

Note that $a_i^2 \neq 0$, unless $b_i \varphi_i(N_i) b_i \varphi_i(N_i) = 0$. For example,if $\varphi_i(N_i) = \frac{1-(-)^{N_i}}{2}$, then $a_i^2 = 0$, implying the hard-core condition for the i^{th} oscillator. Generally , there are other mappings of Bose algebra, equations (1) and (2), but, in general, they do not have the number operators $N_i^{(a)}$, and equation (4) does not hold for other mappings than those in equation (3).

We point out that the complete deformed algebra is associative owing to the mapping of Bose algebra. The Fock space for the deformed algebra is spanned by powers of the creation operators a_i^+,$i \in S$, acting on the vacuum $|0>^{(a)} = |0>^{(b)} \equiv |0>$. The states in the Fock space are specified by the eigenvalues of the number operators N_i, namely $|n_1, n_2, ...n_i.... >^{(a)} = |n_1, n_2, ...n_i.... >^{(b)}$.(If there exists a number $n_i^{(0)} \in \mathbf{N}$, such that $\varphi_i(n_i^{(0)}) = 0$, then $N_i = 0, 1, ...(n_i^{(0)} - 1)$).
States with unit norm are

$$\begin{gathered} |n_1.....n_n >= \frac{(a_1^+)^{n_1}.......(a_n^+)^{n_n}}{\sqrt{[\tilde{\varphi}_1(n_1)]!.....[\tilde{\varphi}_n(n_n)]!}} e^{-\frac{1}{2}\sum_j \theta_{ji}(c_{ij}+c_{ij}^*)n_i n_j} |0, 0, ...0> \\ [\tilde{\varphi}(n)]! = \tilde{\varphi}(n)\tilde{\varphi}(n-1)......\tilde{\varphi}(1) \\ \tilde{\varphi}_i(n_i) = |\varphi_i(n_i)| e^{(c_{ii}+c_{ii}^*)n_i}. \end{gathered} \tag{6}$$

where θ_{ij} is the step function. (For anyons in $(2+1)$ dimension [8], θ is the angle function.)
Furthermore, the matrix elements of the operators a_i, a_i^+, $i \in S$, are

$$\begin{gathered} < ....(n_i - 1)....|a_i|....n_i... >=< ...n_i....|a_i^+|....(n_i - 1).... >^* \\ = \sqrt{\varphi_i(n_i)} e^{\frac{1}{2}\sum_j (c_{ij}+c_{ij}^*)n_j} \prod_{j \neq i} \delta_{n_j, n_j'}. \end{gathered} \tag{7}$$

We also find for any $k = 0, 1, 2, ...$ that

$$\begin{gathered} (a_j^+)^k (a_j)^k = \frac{[\tilde{\varphi}_j(N_j)]!}{[\tilde{\varphi}_j(N_j - k)]!} e^{k\sum_{l \neq j}(c_{jl}+c_{jl}^*)N_l} \\ (a_j)^k (a_j^+)^k = \frac{[\tilde{\varphi}_j(N_j + k)]!}{[\tilde{\varphi}_j(N_j)]!} e^{k\sum_{l \neq j}(c_{jl}+c_{jl}^*)N_l} \end{gathered} \tag{8}$$

The norms of arbitrary linear combinations of the states in equation (6) in the Fock space corresponding to deformed algebra, are positive definite owing to the mapping

of Bose algebra, equations (1) and (2). Namely, $|n_1, n_2, ...n_i.... >^{(a)}=$
$= |n_1, n_2, ...n_i.... >^{(b)} \equiv |n_1, n_2, ...n_i.... >$.

This class of deformed multimode oscillator algebras comprises multimode Biedenharn-Macfarlaine [3], Aric-Coon [6], two-(p,q) parameter [11], Fermi, generalized Green's [12][13], as well as anyonic [8] and Pusz-Woronowicz (with or without the hard-core condition) oscillators covariant under the $SU_q(n)$ ($SU_q(n|m)$) algebra (superalgebra) [7].

Non-isomorphic (non-equivalent) algebras are classified by different matrix elements given by (7), i.e. with the functions $g_i(n_1, n_2, ...n_n) = |\varphi_i(n_i)|e^{\sum_j (c_{ij}+c^*_{ij})n_j}$. It is important to mention that there are mappings of Bose algebra which do not preserve the relation $N_i^{(a)} = N_i^{(b)}$, given by (4). Moreover, there are mappings of Bose algebra for which the number operators $N_i^{(a)}$do not even exist. Such an example is the exchange algebra presented in [14].

Finally, we construct and discuss the number operators $N_i^{(a)}$.Let $\varphi_i(n_i)$ be bijective mappings. Then, using equation (3) we have

$$\begin{gathered} b_i = a_i e^{-\sum_j c_{ij}N_j}\sqrt{\frac{N_i}{\varphi_i(n_i)}}, \quad \forall i \in S \\ N_i = b_i^+ b_i. \end{gathered} \tag{9}$$

The spectra of $N_i^{(a)}$ and $N_i^{(b)}$ coincide. In this case,

$$a_i^+ a_i = g_i(N_1, N_2, ...N_i...) = |\tilde{\varphi}_i(N_i)|e^{\sum_{j\neq i}(c_{ij}+c^*_{ij})N_j}. \tag{10}$$

Let us denote $a_i^+ a_i = x_i$, and remark that x_i commute with any N_j and among themselves. Then the N_i operators can be written as

$$\begin{gathered} N_i = N_i(x_1, ...x_n) = \textstyle\sum_{k=0}^{\infty} \sum_{(i_1...i_k)} \frac{1}{k!}(\frac{\partial^k N_i}{\partial x_{i_1}..\partial x_{i_k}})_{x=0}(x_{i_1}x_{i_2}.....x_{i_k}) \\ N_i(0, 0, ..0) = 0, \quad \forall i \in S. \end{gathered} \tag{11}$$

The coefficients in the Taylor expansion can be obtained from (10)

$$g_i(N_1, N_2, ...N_i...) = x_i, \quad \forall i \in S \tag{12}$$

namely from

$$(\frac{\partial^k g_i(N_1, N_2, ...N_i...)}{\partial x_{j_1}..\partial x_{j_k}})_{x=0} = \delta_{k1}\delta_{ji}. \tag{13}$$

These equations give a set of recurrence relations for the coefficients in the Taylor expansion of N_i as a function of the variables x_j, $j \in S$. For example,

$$\begin{gathered} c_0^{(i)} \equiv (N_i)_{x=0} = 0 \\ c_j^{(i)} \equiv (\tfrac{\partial N_i}{\partial x_j})_{x=0} = \frac{1}{(\frac{\partial g_i}{\partial N_i})_0}\delta_{ij} \\ c_{jk}^{(i)} \equiv \tfrac{1}{2}(\tfrac{\partial^2 N_i}{\partial x_j \partial x_k})_{x=0} = -\tfrac{1}{2}(\tfrac{\partial^2 g_i}{\partial N_j \partial\ N_k})_0 c_i^{(i)} c_j^{(j)} c_k^{(k)}. \end{gathered} \tag{14}$$

Specially

$$\begin{gathered} c_j^{(i)} = \frac{1}{(\frac{d\varphi_i}{dN_i})_0}\delta_{ij} \\ c_{ij}^{(i)} = -\tfrac{1}{2}(c_{ij} + c_{ji}^*)c_i^{(i)} c_j^{(j)} \\ c_{jk}^{(i)} = 0, \quad j \neq i, \quad k \neq i \end{gathered} \tag{15}$$

where $g_i(N_1, N_2, ...N_i...)$ is given by equation (10). We also mention the following result.

If there exists a continuous mapping defined by (3) from Bose oscillators to deformed oscillators, then the number operators exist and can be written in the form

$$N_i = x_i[c_i^{(i)} + \sum_j c_j^{(i)} x_j + ...] \tag{16}$$

where $x_j = a_j^+ a_j$.

For example, let us consider the n-mode Pusz-Woronowicz oscillator algebra[2] (of Bose type, covariant under the $SU_q(n)$ quantum algebra), $q \in \mathbf{R}$:

$$\begin{aligned} a_i a_j &= q^{sgn(j-i)} a_j a_i \\ a_i a_j^+ &= q a_j^+ a_i, \quad i \neq j \\ a_i a_i^+ &= 1 + (q^2-1) \textstyle\sum_j \theta_{ij} a_j^+ a_j + q^2 a_i^+ a_i. \end{aligned} \tag{17}$$

There exist mappings to the Bose algebra (1) and (2) with $c_{ij} = \theta_{ij} \ln q$, $\varphi(N) = \frac{q^{2N}-1}{q^2-1}$ (see Table 1). Hence, we can calculate the coefficients in the expansion (16). Using equation (15) we obtain

$$\begin{aligned} c_j^{(i)} &= \tfrac{q^2-1}{\ln q^2} \delta_{ij} \\ c_{ij}^{(i)} &= -\tfrac{1}{2} \tfrac{(q^2-1)^2}{\ln q^2} (\delta_{ij} + \theta_{ij}). \end{aligned} \tag{18}$$

When $q^2 \to 1$, then $c_i^{(i)} \to 1$, whereas $c_{ij}^{(i)} \to 0$, etc., and $N_i \to a_i^+ a_i$. However, when $q^2 \to 0$, all coefficients diverge and the expansion (16) is not valid. Note that when $q = 0$, the mapping (9) becomes singular, but one can still define the corresponding number operators. For $q = 0$, the Pusz-Woronowicz algebra (17) reduces to

$$\begin{aligned} a_i a_j &= 0, \quad i < j \\ a_i a_j^+ &= 0, \quad i \neq j \\ a_i a_i^+ &= 1 - \textstyle\sum_j \theta_{ij} a_j^+ a_j. \end{aligned} \tag{19}$$

Therefore we also present the number operators in another form that holds for an arbitrary algebra having number operators, and holds even for $q = 0$[12]:

$$N_i = a_i^+ a_i + \sum_{k=1}^{\infty} \sum_{\pi \in S_k} \sum_{j_1 \ldots j_k} d_{\pi(j_1 \ldots j_k), j_1 \ldots j_k} a_{\pi(j_k)}^+ \ldots\ldots a_{\pi(j_1)}^+ a_i^+ a_i a_{j_1} \ldots a_{j_k} \tag{20}$$

where S_k denotes the permutation group and $d_{\pi(j_1 \ldots j_k), j_1 \ldots j_k}$ are the coefficients of the expansion.

For the Pusz-Woronowicz algebra the number operators in the above form (see also [7]) are given by

$$N_i = \sum_{k=0}^{\infty} \sum_{j=0}^{k} c_{k,j} \sum_{p_1 \ldots p_j} \theta_{ip_1} \ldots.. \theta_{ip_j} a_{p_j}^{+} \ldots.. a_{p_1}^{+} (a_i^{+})^{k-j+1} (a_i)^{k-j+1} a_{p_1} \ldots\ldots a_{p_j} \tag{21}$$

with the conditions $c_{0,0} = 1$, $c_{0,-j} = 0$, $j > 0$, and with the recurrence relations

$$c_{k+1,k+1-j} = \frac{(1-q^2)(1-q^{2j})}{1-q^{2(j+1)}} c_{k,k+1-j} + q^{2j}(1-q^2) c_{k,k-j}. \tag{22}$$

Starting with $c_{0,0} = 1$, we find that

$$\begin{aligned} c_{k,1} = c_{k,k} &= (1-q^2)^k \\ c_{k,0} &= \tfrac{(1-q^2)^{k+1}}{1-q^{2(k+1)}}. \end{aligned} \tag{23}$$

In the limit $q^2 \to 1$, the number operator is $N_i = a_i^{+} a_i$. In the limit $q = 0$, all coefficients are $c_{k,j} = 1$, $\forall k, j$. This result is similar to that found by Greenberg [9] for quons with $q = 0$. Finally, for $\varphi_i(N_i) = \frac{1-(-)^{N_i}}{2}$, the number operators, $N_i^{(a)} \neq N_i^{(b)}$, are simply

$$N_i^{(a)} = a_i^{+} a_i = b_i^{+} b_i [1 - \theta(n_i - 1)]. \tag{24}$$

We conclude that we unify and generalize the results for the states in the Fock space and the number operators for the mappings of multimode Bose algebra presented in Table 1.

Acknowledgments

This work was supported by the joint Croatian-American contract NSF JF 999 and the Scientific Fund of the Republic of Croatia.

References

[1] Drinfeld V G 1986 *in Proceedings of the ICM (Berkeley, CA)* p.798
Jimbo M 1985 *Lett.Math.Phys.* **10** 63

[2] Pusz W and Woronowicz S L 1989 *Rep.Math.Phys.* **27** 231

[3] Biedenharn L C 1989 *J. Phys.A :Math.Gen.* **22** L873
Macfarlane A J 1989 *J. Phys.A :Math.Gen.* **22** L983

[4] Hayashi T 1990 *Comm.Math.Phys.* **127** 129

[5] Odaka K, Kishi T and Kamefuchi S 1991 *J. Phys.A :Math.Gen.* **24** L591
Chaturvedi S and Srinivasan V 1992 *Phys.Rev.A* **44** 8020

[6] Arik M and Coon D D 1976 *J.Math.Phys.* **17** 524

[7] Jagannathan R et al. 1992 *J. Phys.A :Math.Gen.* **25** 6429

[8] Lerda A and Sciuto S 1993 *Nucl.Phys.B* **401** 613
Bardek V, DorešićM and Meljanac S 1993 Example of q-deformed field theory, hep-th 9310171 *(to appear in Int.J.Mod.Phys)*
Bardek V, DorešićM and Meljanac S 1993 Anyons as quons, *(to appear in Phys.Rev.D)*

[9] Greenberg O W 1990 *Phys.Rev.Lett.* **64** 705
Greenberg O W 1991*Phys.Rev.D* **43** 4111

[10] Fairlie D and Zachos C 1991 *Phys.Lett.* **256B** 43
Van der Jeugt J 1993 *J. Phys.A :Math.Gen.* **26** L405

Meljanac S, MilekovićM and Perica A 1993 On the R-matrix formulation of deformed algebras *(submitted to J. Phys.A :Math.Gen.)*

[11] Chakrabarti R and Jagannathan R 1991 *J. Phys.A :Math.Gen.* **24** L711
Kibler M R 1992 Introduction to quantum algebras *preprint LYCEN 9234*

[12] Meljanac S and Perica A 1993 Bose, Fermi, para-Bose, para-Fermi and anyon statistics: The R- matrix approach to deformed associative algebras, *(submitted to Phys.Rev.Lett.)*

[13] Ohnuki Y and Kamefuchi S, 1982 *Quantum Field Theory and Parastatistic (University of Tokio Press, Tokio, Springer, Berlin)*

[14] Brink L, Hansson T H, Konstein S and Vasiliev M A 1993 *Nucl.Phys.* **401B** 591
Polychronakos A P 1992 *Phys.Rev.Lett.* **69** 703

Table 1: Parameters c_{ij} and $\varphi_i(N_i)$ for the various deformed algebras.We use definition $[x]_p = \frac{p^x-1}{p-1}$.

type of oscillators	c_{ij}	$\varphi_i(N_i)$
Bose	$i\pi\lambda_{ij}$, $\lambda_{ij} - \lambda_{ji} \in 2\mathbf{Z}$, $\forall i, j$	N_i
Fermi	$i\pi\lambda_{ij}$, $\lambda_{ij} - \lambda_{ji} \in 2\mathbf{Z} + 1$, $\forall i, j$	$[N_i]_{-1}$
Green's oscillators [12][13]	$i\pi\lambda_{ij}$, $\lambda_{ij} - \lambda_{ji} \in \mathbf{Z}$, $\forall i, j$	$[N_i]_{\pm 1}$
Anyonic-type [10]	$i\pi\lambda_{ij}$, $\lambda_{ij} - \lambda_{ji} \in \mathbf{R}$, $\forall i, j$	$[N_i]_{\pm 1}$
Anyons [8]	$i\lambda\theta_{ij}$, $\lambda \in \mathbf{R}$, θ_{ij} is angle	$[N_i]_{\pm 1}$
Pusz-Woronowicz (Bose) [2]	$\theta_{ij} \ln q$, $q \in \mathbf{R}$	$[N_i]_{q^2}$
Pusz-Woronowicz (Fermi) [2]	$\theta_{ij} \ln(-q)$, $q \in \mathbf{R}$	$[N_i]_{-1}$
$SU_q(n\|m)$-covariant (Bose) oscillators [7]	$\theta_{ij} \ln q$, $q \in \mathbf{R}$, $i \leq n$	$[N_i]_{q^2}$
$SU_q(n\|m)$-covariant (Fermi) oscillators [7]	$\theta_{ij} \ln(-q)$, $q \in \mathbf{R}$, $n + 1 \leq i$	$[N_i]_{-1}$
Biedenharn-Macfarlane [3]	$-\frac{1}{2}\delta_{ij} \ln q_i$, $q_i \in \mathbf{C}$	$[N_i]_{q_i^2}$
Arik-Coon [6]	0, $q_i \in \mathbf{C}$	$[N_i]_{q_i}$